




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>G01N 33/558, C12Q 1/68</b>		<b>A1</b>	(11) International Publication Number: <b>WO 95/17676</b>
			(43) International Publication Date: 29 June 1995 (29.06.95)
(21) International Application Number: PCT/EP94/04232 (22) International Filing Date: 20 December 1994 (20.12.94) (30) Priority Data: 108159 23 December 1993 (23.12.93) IL (71) Applicant (for all designated States except US): ORGENICS INTERNATIONAL HOLDINGS B.V. [NL/NL]; c/o TMF Management B.V., Emmamplein 5, NL-1075 AW Amsterdam (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): RITTERBAND, Menachem [IL/IL]; 9B Maccabi Street, 70400 Ness Ziona (IL). ALAJEM, Sarah [IL/IL]; 64 Kfar Hanagid 76875 (IL). SCHRIFT, Michal [IL/IL]; 6 Hapalmach Street, 82000 Kiryat Gat (IL). RUSU, Ileana [IL/IL]; 70/3 Derech Yavne, 76341 Rehovot (IL). REINHARTZ, Avraham [IL/IL]; 1 Israel Shahr Street, 76450 Rehovot (IL). (74) Agents: BAILLIE, Iain, C. et al.; Ladas & Parry, Altheimer Eck 2, D-80331 München (DE).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published With international search report.	

(54) Title: APPARATUS FOR SEPARATION, CONCENTRATION AND DETECTION OF TARGET MOLECULES IN A LIQUID SAMPLE

## (57) Abstract

This disclosure describes an apparatus (10) for separation and detection of target molecules in a liquid sample containing the target molecules and non-target molecules comprising at least one absorption area (20), a bibulous carrier (18) defining a first liquid transport path in fluid contact with at least one absorption area (20) including at least one reaction area (26, 30) upstream of the at least one absorption area (20) wherein the reaction area (26, 30) includes a capture reagent and a sample area (24) upstream of the reaction area (26, 30) wherein the liquid sample when applied to sample area (28) is transported by capillary action past the reaction area (26, 30) where at least a portion of the target molecules may be bound to the capture reagent to permit detection of the target molecules. The liquid sample is then transported by capillary action on to the absorption area (20). A second liquid transport path in fluid communication with the reaction area (26, 30) of the bibulous carrier (18) for transporting a washing solution applied to the reaction area (26, 30) to the at least one absorption area (20) thereby removing interfering substances from the reaction area (26, 30). Means are also provided for alternatively selecting between the first and second liquid transport path during capillary transport of the liquid sample.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

APPARATUS FOR SEPARATION, CONCENTRATION AND DETECTION OF  
TARGET MOLECULES IN A LIQUID SAMPLE

The invention relates to apparatus for separation concentration and detection of target molecules in a liquid sample.

The use of test devices including chromatography as a separation, concentration and detection procedure is well known in the art. For example, European Patent Application 0 262 328 A3 describes a test strip for the detection of analytes such as antigens and antibodies. Chromatographic test devices, such as the device described in U.S. Patent 4,857,453 to Ullman et al, may also be enclosed in a housing.

One limitation of chromatography devices such as those described above is that washing procedures are slow since the washing fluid must pass through the entire length of the chromatographic path. A device in which a washing solution is carried only over a portion of the chromatographic path is described in PCT Application WO 91/15769.

The present invention provides an improved apparatus for separation, concentration and detection of target molecules in a liquid sample.

There is thus provided apparatus for separation, concentration and detection of target molecules in a liquid sample containing the target molecules and non-target molecules comprising, at least one absorption area, a bibulous carrier defining a first liquid trans-

port path in fluid contact with the at least one absorption area including, at least one reaction area upstream of the at least one absorption area wherein the reaction area includes a capture reagent, and a sample area upstream of the reaction area wherein the liquid sample when applied to sample area is transported by capillary action past the reaction area where at least a portion of the target molecules may be bound to the capture reagent to permit detection of the target molecules and the liquid sample is then transported by capillary action on to the absorption area, a second liquid transport path in fluid communication with the reaction area of the bibulous carrier for transporting a washing solution applied to the reaction area to the at least one absorption thereby removing interfering substances from the reaction area, and apparatus for alternatively selecting between the first and second liquid transport path during capillary transport of the liquid sample.

In accordance with a preferred embodiment of the present invention the apparatus for separation concentration and detection is enclosed in a housing.

In accordance with a another preferred embodiment of the present invention the housing includes a sample port in fluid communication with the sample area.

In accordance with yet another preferred embodiment of the present invention the housing includes a washing port in fluid communication with the reaction area.

In accordance with still another preferred embodiment of the present invention the at least one absorption area includes a first absorption area and a second absorption area.

In accordance with yet another preferred embodiment of the present invention the second absorption area is in direct fluid connection with the first absorption area for prevention of a backflow from the first absorption area through bibulous carrier to the second absorption

area.

In accordance with a further preferred embodiment of the present invention the bibulous carrier is a nitrocellulose membrane wherein the absorption sites have been blocked to facilitate transport of the target molecules.

In accordance with a still further preferred embodiment of the present invention the bibulous carrier is a glass fiber matrix.

In accordance with another preferred embodiment of the present invention the apparatus for alternatively selecting includes a fluid impermeable disintegratable partition between the bibulous carrier and the second absorption area.

In accordance with yet another preferred embodiment of the present invention the disintegratable partition is disintegrated by the liquid sample.

In accordance with still another preferred embodiment of the present invention the disintegratable partition is disintegrated by a substance added to the liquid sample.

In accordance with yet another preferred embodiment of the present invention the disintegratable partition is disintegrated by a solution applied to the sample area after the liquid sample is applied.

In accordance with a further preferred embodiment of the present invention the disintegratable partition is disintegrated by a substance applied to the reaction area after binding of the target molecules to the reaction area.

In accordance with a still further preferred embodiment of the present invention the disintegratable partition is disintegrated over a predetermined period of time.

In accordance with yet a further preferred embodiment of the present invention the disintegratable partition is fabricated from a netting embedded in a compound

selected from the group including water soluble polymers, pH degradable substances, ionic strength degradable substances and enzymatically digestible substances.

In accordance with another preferred embodiment of the present invention the water soluble polymers include water soluble polyesters.

In accordance with yet another preferred embodiment of the present invention the target molecules include target nucleic acid sequences.

In accordance with still another preferred embodiment of the present invention the target nucleic acid sequences include DNA sequences.

In accordance with yet another preferred embodiment of the present invention the target nucleic acid sequences include RNA sequences.

In accordance with a further preferred embodiment of the present invention the target molecules include antigens.

In accordance with still a further preferred embodiment of the present invention the target molecules include antibodies.

In accordance with yet a further preferred embodiment of the present invention the capture reagent comprises at least one nucleic acid capture reagent including nucleic acid probe sequences complementary to at least part of the target nucleic acid sequences.

In accordance with another preferred embodiment of the present invention the capture reagent comprises at least one antigen.

In accordance with yet another preferred embodiment of the present invention the capture reagent comprises at least one antibody.

The present invention will be understood and appreciated more fully from the following detailed description taken in conjunction with the drawings in which:

Fig. 1 is an exploded pictorial illustration of apparatus for separation, concentration and detection of target molecules in a liquid sample constructed and operative in accordance with the present invention; and

Fig. 2 is a side sectional illustration of the apparatus taken along the line II - II of Fig. 1.

Reference is now made to Figs. 1 and 2 which illustrate apparatus 10 for separation, concentration and detection of target molecules in a liquid sample constructed and operative in accordance with a preferred embodiment of the present invention.

Apparatus 10 includes a housing 12 fabricated from an injection moldable, radiation sterilizable plastic such as polyethylene. The housing 12 includes a top member 14, and a bottom member 16 which are joined together by any convenient means such as gluing.

The housing typically encloses a bibulous matrix 18, an absorption area 20 and a disintegratable barrier 22. The bibulous matrix 18 when wet forms a capillary transport path in fluid communication with the absorption area and includes a sample area 24 upstream of a reaction area 26 which is in fluid communication with the absorption area 20.

Typically, a liquid sample of 0.1 ml to 1.0 ml in volume including target and non-target molecules is applied to the sample area 24 through a sample port 28 in the top member 14 of the housing 12. The liquid sample is then carried along a first transport path in the bibulous matrix 18 by capillary action from the sample area 24 to the reaction area 26 where the target molecules may be captured by a capture reagent for later detection.

The capture reagent is typically an antibody to the target molecule or a DNA or RNA sequence complementary to at least a portion of the target molecule. Detection is typically accomplished by a colorimetric reaction in the reaction area 26 between a detection reagent and the captured target molecules.

From the reaction area 26 the liquid sample is typically carried along the first transport path to the absorption area 20 by capillary action. A portion 31 of

the absorption area 20 typically abuts a lower surface of the bibulous matrix 18 downstream of the reaction area 26 to maximize capillary flow forces in the bibulous matrix 18.

The bibulous matrix 18 and the absorption area 20 are typically fixed to the top member 14 by a pressure sensitive, low soak in adhesive such as depositable adhesive, I.D. No. 62-9985-4830-4, 3M Industrial Specialties Div., St. Paul MN, USA.

The disintegratable barrier 22 is typically disposed between the first flow path and the absorption area 20 upstream of portion 31. The disintegratable barrier 22 typically abuts and separates the absorption area 20 from the bibulous matrix 18 from a point downstream of the sample port 28 to part of the absorption area 20 downstream of reaction area 26 and upstream of portion 31 of the absorption area 20. The disintegratable barrier 22 typically extends laterally beyond the bibulous matrix 18 and the absorption area 20 and is typically fixed to the top member 14 by a pressure sensitive, low soak in adhesive in the areas not covered by the bibulous matrix 18 or the absorption area 20. The lateral extension of the disintegratable barrier 22 beyond the absorption area 20 blocks direct fluid contact between the bibulous matrix 18 and the absorption area 20 except at portion 31, until the disintegratable barrier 22 is disintegrated.

When the disintegratable barrier 22 is disintegrated a second transport path is established for capillary transport between the reaction area 26 and the absorption area 20 through the bibulous matrix 18. This second transport path can be employed to wash interfering compounds from the reaction area 26 by applying a washing solution to the reaction area 26 through wash port 30 in top member 14.

The disintegratable barrier 22 may be typically

fabricated from compounds embedded on a netting platform of plastic polymeric material. These compounds typically include water soluble polymers such as the polyesters lactic acid and glycolic acid copolymers, pH and/or ionic strength degradable substances such as poly-hydroxymethylacrylate and carbopol and enzymatically digestible substrates such as gelatin which may be digested by gelatinase.

The absorption area 20 is typically fitted into a well 32 in bottom member 16. The absorption area 20 is typically fabricated to absorb all of the liquid to be added to apparatus 10 during use in an assay procedure.

The procedure used for the separation, concentration and detection of target molecules using apparatus 10 typically includes applying 0.05 ml to 0.5 ml of a sample fluid to the sample area 24 of the bibulous matrix 18 through sample port 28. A wash fluid may subsequently be applied to the sample area 24 when the sample fluid volume is insufficient for efficient capillary transport of the sample liquid to the reaction area 26. Either the sample fluid or the wash fluid may contain a compound which disintegrates the disintegratable barrier 22.

Capillary and chromatographic forces within the bibulous matrix 18 draw the fluid portion of the sample along the first transport path from the sample area 24 toward the reaction area 26 and on to the absorption area 20. As the fluid migrates through the reaction area 26, target molecules in the sample fluid react and bind with a specific complementary capture reagent such as an immuno-reagent, a chemical test or a nucleic acid sequence reagents, which has been immobilized to the reaction area 26.

The absorptive capacity of portion 31 and contact between the absorption area 20 and the bibulous matrix 18 at portion 31 prevents a counter flow of fluid along the first transport path from the absorption area 20 back to

the reaction area 26 if the bibulous matrix 18 is overloaded. Thus any excess fluid applied to the bibulous matrix 18 or the absorption area 20 will flow into the portion 31 rather than counter flow.

As the fluids are transported in the bibulous matrix 18 along the first transport path, the disintegrative compound contained in either the sample or wash fluid start to cleave the disintegratable barrier 22 which separates the bibulous matrix 18 and the absorption area 20. The time elapsed from the application of the disintegrative compound to disintegration of the disintegratable barrier 22 may typically range from 0.5 to 45 minutes depending on the composition of the disintegratable barrier.

After capillary transport of the fluids applied to the sample area 24 is completed a suitable volume of wash fluid, typically 0.1 ml to 1.5 ml, may be applied directly to the reaction area 26 through the reaction port 30. Typically, the wash fluid applied to the reaction area 26 completes the cleavage of the disintegratable barrier 22 thereby establishing the second transport path through the bibulous matrix 18 between the reaction area 26 and the absorption area 20. This second flow path permits the rapid and efficient washing of the reaction area 26 to remove any unreacted sample molecules or other compounds which may interfere with subsequent steps. The wash fluid applied directly to the reaction area 26 may, however, alternatively contain the disintegrative compound.

After washing of the reaction area 26 an immuno-reagent, or chemical test reagent complementary to the target molecules conjugated with an enzyme or other suitable tracer, is applied directly to the reaction area 26 through reaction port 30. Most of the unbound immuno-reagent conjugate or chemical test reagent is then washed out of the reaction area 26 into the absorption area 20 along the second flow path by application of a washing

fluid to the reaction area 26. If the tracer is an enzyme requiring a substrate to develop color, a volume of this substrate or chromogen is applied to the reaction area 26 through reaction port 30 immediately after washing of unbound reagent.

Reference is now made to the following examples which, together with Figs. 1 and 2 illustrate the invention.

#### EXAMPLE 1

##### DETECTION OF HIV

Synthetic HIV - 1 (gp - 41, gp -120) antigen peptides was immobilized on the reaction zone of a glass fiber matrix. Approximately 300 microliters of pre-diluted blood serum was applied to the sample area 28 of apparatus 10. This was followed by application to the sample area 28 of 150 microliters of a wash solution consisting of 1 % gelatin and 0.3 % tween-20 in a phosphate buffered saline.

Immediately after the transport of fluids to the reaction areas 30 as indicated by the wetting of the reaction area 30 and the disappearance of liquid from the sample area 28, 300 microliters of an affinity purified rabbit anti-human IgG, conjugated to alkaline phosphatase, diluted in a conjugate solution of greater viscosity comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, was directly applied to the wetted reaction zone. Unreacted enzyme conjugate was washed out of the reaction area 30 by applying 500 microliters of wash solution to the reaction area 30. Finally, 100 microliters of a substrate chromogen BCIP/NBT was applied to the reaction area 30. Appearance of a colored product at the reaction zone provided evidence of enzyme activity indicative of antibody to HIV -1 present in the sample.

EXAMPLE 2  
DETECTION OF HIV-1

Recombinant HIV - 1 (gp - 41) antigen was immobilized on the reaction zone of a glass fiber matrix. After immobilization of the antigen, the reaction zone was blocked by adding 150 ml of matrix blocker comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.05% thimerosal, to the reaction zone area. The matrix blocker migrated by capillary forces and wetted almost the whole glass fiber matrix.

Approximately 300 microliters of pre-diluted blood serum was applied to the sample area 28 of apparatus 10. This was followed by application to the sample area 28 of 150 microliters of a wash solution consisting of 1 % gelatin, 0.05% thimerosal and 0.3 % tween-20 in a phosphate buffered saline.

After the transport of fluids to the reaction areas 30 as indicated by the wetting of the reaction area 30 and the disappearance of liquid from the sample area 28, 300 microliters of an affinity purified rabbit anti-human IgG, conjugated to alkaline phosphatase, diluted in a conjugate solution of greater viscosity comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, was directly applied to the wetted reaction zone. Unreacted enzyme conjugate was washed out of the reaction area 30 by applying 600 microliters of wash solution to the reaction area 30. Finally, 300 microliters of a substrate chromogen BCIP/NBT was applied to the reaction area 30. Appearance of a colored product at the reaction zone provided evidence of enzyme activity indicative of antibody to HIV -1 present in the sample.

EXAMPLE 3  
DETECTION OF HIV-1 IN SALIVA

Synthetic HIV - 1 (gp - 41, gp - 120) antigen peptides was immobilized on the reaction zone of a glass fiber matrix. After immobilization of the antigen, the reaction zone was blocked by adding 150 ul of matrix blocker comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, to the reaction zone area. The matrix blocker migrated by capillary forces and wetted almost the whole glass fiber matrix.

Approximately 200 microliters of pre-treated saliva was then applied to the sample area 28 of apparatus 10. The saliva was pretreated according to the user manual of Omni-Sal<sup>tm</sup>, Saliva Diagnostic Systems Inc., Trantdale, OR, USA.

Immediately after the transport of fluids to the reaction areas 30 as indicated by the wetting of the reaction area 30 and the disappearance of liquid from the sample area 28, approximately 200 microliters of an affinity purified rabbit anti-human IgG, conjugated to Carbon Sol particles, Holland Biotechnology Co., Leiden, the Netherlands, diluted in a conjugate solution of greater viscosity comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, was directly applied to the wetted reaction zone. Unreacted conjugate was washed out of the reaction area 30 by three applications of 500 microliters of wash solution to the reaction area 30. Appearance of a colored spot at the reaction zone provided evidence indicative of antibody to HIV -1 present in the sample.

EXAMPLE 4  
DETECTION OF HELICOBACTER PYLORI

Helicobacter Pylori antigen was immobilized on the reaction zone of a glass fiber matrix. After immobilization of the antigen, the reaction zone was blocked by adding 150 ul of matrix blocker comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, to the reaction zone area. The matrix blocker migrated by capillary forces and wetted almost the whole glass fiber matrix.

Approximately 300 microliters of blood serum pre-diluted in a running solution of 1 % gelatin, 0.05% thimerosal and 0.3 % tween-20 in a phosphate buffered saline was applied to the sample area 28 of apparatus 10.

Immediately after the transport of fluids to the reaction areas 30 as indicated by the wetting of the reaction area 30 and the disappearance of liquid from the sample area 28, 300 microliters of an affinity purified rabbit anti-human IgG, conjugated to alkaline phosphatase, diluted in a conjugate solution of greater viscosity comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, was directly applied to the wetted reaction zone. Unreacted enzyme conjugate was washed out of the reaction area 30 by applying 600 microliters of wash solution to the reaction area 30. Finally, 300 microliters of a substrate chromogen BCIP/NBT was applied to the reaction area 30. Appearance of a colored product at the reaction zone provided evidence of enzyme activity indicative of antibody to Helicobacter Pylori present in the sample.

EXAMPLE 5  
DETECTION OF HCV

NS-4a-peptide, NS-3 recombinant protein and the core structural peptide of the HCV antigen were immobilized on the reaction zone of a glass fiber matrix. After immobilization of the antigen, the reaction zone was blocked by adding 150 ul of matrix blocker comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, to the reaction zone area. The matrix blocker migrated by capillary forces and wetted almost the whole glass fiber matrix.

Approximately 300 microliters of blood serum pre-diluted in a running solution of 1 % gelatin, 0.05% thimerosal and 0.3 % tween-20 in a phosphate buffered saline was applied to the sample area 28 of apparatus 10.

Immediately after the transport of fluids to the reaction areas 30 as indicated by the wetting of the reaction area 30 and the disappearance of liquid from the sample area 28, 300 microliters of an affinity purified rabbit anti-human IgG, conjugated to alkaline phosphatase, diluted in a conjugate solution of greater viscosity comprising a phosphate buffered saline solution including 1% gelatin, 1% fetal calf serum, 0.05% tween 20, 1% PVP 700,000 and 0.5% thimerosal, was directly applied to the wetted reaction zone. Unreacted enzyme conjugate was washed out of the reaction area 30 by applying 600 microliters of wash solution to the reaction area 30. Finally, 300 microliters of a substrate chromogen BCIP/NBT was applied to the reaction area 30. Appearance of a colored product at the reaction zone provided evidence of enzyme activity indicative of antibody to HCV present in the sample.

EXAMPLE 6  
DETECTION OF HB<sub>s</sub> ANTIGEN

A mixture of two different monoclonal anti HB<sub>s</sub> antibodies in a phosphate buffered saline solution including 0.005% bovine serum albumin and 2% trehalose were immobilized on the reaction zone of a glass fiber matrix. After immobilization of the antigen, the reaction zone was blocked by adding 150 ul of matrix blocker comprising a phosphate buffered saline solution including 2% bovine serum and 1% PVP 700,000, to the sample area 28. The matrix blocker migrated by capillary forces and wetted almost the whole glass fiber matrix.

Approximately 200 microliters of blood serum pre-diluted in a phosphate buffered saline reaction solution of greater viscosity consisting of 35% serum sample, 0.5% PVP 700000, 1.0% fetal calf serum, 0.002% horse serum, 4.0% dextran sulfate, 0.04% tween 20 and precalibrated volumes of biotinylated monoclonal anti HB<sub>s</sub> antibodies and strepavidin APL conjugate was then directly applied to the sample area 28 of apparatus 10.

Immediately after the transport of fluids to the reaction area 30 as indicated by the wetting of the reaction area 30 and the disappearance of liquid from the sample area 28 unreacted enzyme conjugate was washed out of the reaction area 30 by four applications of 500 microliters of wash solution to the reaction area 30. Finally, 200 microliters of a substrate chromogen BCIP/NBT was applied to the reaction area 30. Appearance of a colored product at the reaction zone provided evidence of enzyme activity indicative of HB<sub>s</sub> antigen present in the sample.

EXAMPLE 7  
DETECTION OF HPV BY HYBRIDIZATION

HPV specific oligonucleotides (GTTTCAGGACCCACAGGAGC-GACCC) was immobilized on the reaction zone of a glass fiber matrix. DNA from Caski cells was extracted and labeled through sulfonation according to the protocol of Hybricomb<sup>tm</sup>, Organics Ltd., Yavne, Israel. The labeled DNA was then diluted 1:10 in a hybridization solution composed of 0.6 M NaCl, 20mM phosphate buffer, pH 7.5, 0.02% Ficoll, 0.02% gelatin, 0.1% tween 20 and 20.0% glycerol.

The sample was then boiled for 20 minutes and then immediately chilled on ice. 200 ul of the solution was applied to

the sample area 28 of apparatus 10. After the transport of all the hybridization solution, 100 ul of an affinity purified mouse anti sulfonated DNA conjugated to alkaline-phosphatase diluted in a phosphate buffered saline solution including 1% gelatin and 0.1% tween was then applied to the reaction area 30.

Unreacted enzyme conjugate was washed out of the reaction area 30 by applying 500 microliters of a phosphate buffered saline wash solution including 0.5% tween to the reaction area 30. Finally, 100 microliters of a substrate chromogen BCIP/NBT was applied to the reaction area 30. Appearance of a colored product at the reaction zone provided evidence of enzyme activity indicative specific HPV-16 sequences present in the sample.

It will be appreciated by persons skilled in the art that the present invention is not limited to what has been particularly shown and described herein above. Rather the scope of the present invention is defined only by the claims which follow:

## CLAIMS

1. Apparatus for separation, concentration and detection of target molecules in a liquid sample containing the target molecules and non-target molecules comprising:

at least one absorption area;

a bibulous carrier defining a first liquid transport path in fluid contact with the at least one absorption area including:

at least one reaction area upstream of the at least one absorption area wherein the reaction area includes a capture reagent; and

a sample area upstream of the reaction area wherein the liquid sample when applied to sample area is transported by capillary action past the reaction area where at least a portion of the target molecules may be bound to the capture reagent to permit detection of the target molecules and the liquid sample is then transported by capillary action on to the absorption area;

a second liquid transport path in fluid communication with the reaction area of the bibulous carrier for transporting a washing solution applied to the reaction area to the at least one absorption area thereby removing interfering substances from the reaction area; and

means for alternatively selecting between the first and second liquid transport path during capillary transport of the liquid sample.

2. Apparatus according to claim 1 wherein the apparatus for separation concentration and detection is enclosed in a housing.

3. Apparatus according to claim 2 wherein the housing includes a sample port in fluid communication with the sample area.

4. Apparatus according to claim 2 wherein the housing includes a washing port in fluid communication with the reaction area.

5. Apparatus according to claim 1 wherein the at least one absorption area includes a first absorption area and a second absorption area.

6. Apparatus according to claim 5 wherein the second absorption area is in direct fluid connection with the first absorption area for prevention of a backflow from the first absorption area through bibulous carrier to the second absorption area.

7. Apparatus according to claim 1 wherein the bibulous carrier is a nitrocellulose membrane wherein the absorption sites have been blocked to facilitate transport of the target molecules.

8. Apparatus according to claim 1 wherein the bibulous carrier is a glass fiber matrix.

9. Apparatus according to claim 1 wherein the means for alternatively selecting includes a fluid impermeable disintegratable partition between the bibulous carrier and the second absorption area.

10. Apparatus according to claim 9 wherein the disintegratable partition is disintegrated by the liquid sample.

11. Apparatus according to claim 9 wherein the disintegratable partition is disintegrated by a substance added to the liquid sample.

12. Apparatus according to claim 9 wherein the disintegratable partition is disintegrated by a solution applied

to the sample area after the liquid sample is applied.

13. Apparatus according to claim 9 wherein the disintegratable partition is disintegrated by a substance applied to the reaction area after binding of the target molecules to the reaction area.

14. Apparatus according to claim 9 wherein the disintegratable partition is disintegrated over a predetermined period of time.

15. Apparatus according to claim 9 wherein the disintegratable partition is fabricated from a netting embedded in a compound selected from the group including water soluble polymers, pH degradable substances, ionic strength degradable substances and enzymatically digestible substances.

16. Apparatus according to claim 15 wherein the water soluble polymers include water soluble polyesters.

17. Apparatus according to claim 1 wherein the target molecules include target nucleic acid sequences.

18. Apparatus according to claim 17 wherein the target nucleic acid sequences include DNA sequences.

19. Apparatus according to claim 17 wherein the target nucleic acid sequences include RNA sequences.

20. Apparatus according to claim 1 wherein the target molecules include antigens.

21. Apparatus according to claim 1 wherein the target molecules include antibodies.


22. Apparatus according to claim 17 wherein the capture reagent comprises at least one nucleic acid capture reagent including nucleic acid probe sequences complementary to at least part of the target nucleic acid sequences.

23. Apparatus according to claim 1 wherein the capture reagent comprises at least one antigen.

24. Apparatus according to claim 1 wherein the capture reagent comprises at least one antibody.


1/2

FIG. 1



2/2

FIG. 2



## INTERNATIONAL SEARCH REPORT

PCT/EP 94/04232

## A. CLASSIFICATION OF SUBJECT MATTER

G 01 N 33/558, C 12 Q 1/68

According to International Patent Classification (IPC) or to both national classification and IPC <sup>6</sup>

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

G 01 N, C 12 Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A, 93/07 292 (ORGENICS INTERNATIONAL HOLDINGS B.V.) 15 April 1993 (15.04.93), the whole document, especially claims. --	1, 17- 19, 22
A	WO, A, 91/15 769 (DISEASE DETECTION INTERNATIONAL) 17 October 1991 (17.10.91), abstract; claims; figs.. --	1, 20, 21, 23, 24
A	EP, A, 0 306 336 (SYNTEX (U.S.A.) INC.) 08 March 1989 (08.03.89), abstract; claims; figs.. --	1, 20, 21, 23, 24
A	US, A, 4 857 453	1, 20,

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

09 March 1995

Date of mailing of the international search report

27 March 1995

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

SCHNASS e.h.

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	(ULLMAN et al.) 15 August 1989 (15.08.89), abstract; claims; figs.. --	21, 23, 24
A	EP, A, 0 323 605 (ABBOTT LABORATORIES) 12 July 1989 (12.07.89), abstract; claims; figs.. --	1, 20, 21, 23, 24
A	EP, A, 0 306 772 (ABBOTT LABORATORIES) 15 March 1989 (15.03.89), abstract; claims; figs.. ----	1, 20, 21, 23, 24

## ANHANG

## ANNEX

## ANNEXE

zum internationalen Recherchen-  
bericht über die internationale  
Patentanmeldung Nr.

to the International Search  
Report to the International Patent  
Application No.

au rapport de recherche inter-  
national relatif à la demande de brevet  
international n°

PCT/EP 94/04232 SAE 102312

In diesem Anhang sind die Mitglieder  
der Patentfamilien der im obenge-  
nannten internationalen Recherchenbericht  
angeführten Patentdokumente angegeben.  
Diese Angaben dienen nur zur Unter-  
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family  
members relating to the patent documents  
cited in the above-mentioned inter-  
national search report. The Office is  
in no way liable for these particulars  
which are given merely for the purpose  
of information.

La présente annexe indique les  
membres de la famille de brevets  
relatifs aux documents de brevets cités  
dans le rapport de recherche inter-  
national visée ci-dessus. Les renseigne-  
ments fournis sont donnés à titre indica-  
tif et n'engagent pas la responsabilité  
de l'Office.

In Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
WO A1 9307292	15-04-93	AU A1 27894/92	03-05-93
		EP A1 612354	31-08-94
		IL A0 102486	14-01-93
		IL A0 99647	18-08-92
WO A1 9115769	17-10-91	AU A1 76722/91	30-10-91
EP A2 306336	08-03-89	CA A1 1333046	15-11-94
		DE C0 3887941	31-03-94
		DE T2 3887941	01-09-94
		EP A3 306336	07-11-90
		EP B1 306336	23-02-94
		ES T3 2051303	16-06-94
		JP A2 1072066	16-03-89
		US A 4981786	01-01-91
US A 4857453	15-08-89	AU A1 14305/88	13-10-88
		AU B2 625170	02-07-92
		CA A1 1303491	16-06-92
		DE C0 3851772	17-11-94
		DK A0 1862/88	06-04-88
		DK A 1862/88	08-10-88
		EP A2 286371	12-10-88
		EP A3 286371	07-03-90
		EP A3 286371	07-11-90
		EP B1 286371	12-10-94
		ES T3 2065334	16-02-95
		IL A0 85997	30-09-88
		IL A1 85997	21-06-92
		JP A2 63269056	07-11-88
		NO A0 881480	06-04-88
		NO A 881480	10-10-88
		US A 5137808	11-08-92
EP A2 323605	12-07-89	AT E 100937	15-02-94
		AU A1 27095/88	29-06-89
		AU B2 619231	23-01-92
		CA A1 1312544	12-01-93
		CN A 1036078	04-10-89
		DE C0 3887491	10-03-94
		DE T2 3887491	07-07-94
		EP A3 323605	09-08-89
		EP B1 323605	26-01-94
		ES T3 2050697	01-06-94
		JP A2 1244370	28-09-89
EP A1 306772	15-03-89	AT E 86746	15-03-93
		AU A1 21753/88	16-03-89
		AU B2 619480	30-01-92
		DE C0 3879048	15-04-93
		DE T2 3879048	02-09-93
		EP B1 306772	10-03-93
		ES T3 2039533	01-10-93
		JP A2 1113662	02-05-89
		JP B4 6027738	13-04-94
		US A 4956302	11-09-90

